This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61F 2/04, 2/06, 2/24, 2/54

A1

US

(11) International Publication Number:

WO 97/09006

(43) International Publication Date:

13 March 1997 (13.03.97)

(21) International Application Number:

PCT/US96/13907

(22) International Filing Date:

30 August 1996 (30.08.96)

(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

(30) Priority Data:

08/522,712

1 September 1995 (01.09.95)

Published


With international search report.

- (71) Applicant: EMORY UNIVERSITY [US/US]; 2009 Ridgewood Drive, Atlanta, GA 30322 (US).
- (72) Inventors: WAKSMAN, Ron; 603 Calibre Woods, Atlanta, GA 30329 (US). KING, Spencer, B.; 925 Lullwater Parkway, Atlanta, GA 30307 (US). SCOTT, Neal, A.; 1360 Fairview Road, Atlanta, GA 30306 (US).
- (74) Agents: NEEDLE, William, H. et al.; Needle & Rosenberg, Suite 1200, 127 Peachtree Street, N.E., Atlanta, GA 30303-1811 (US).

(54) Title: ENDOVASCULAR SUPPORT DEVICE AND METHOD OF USE

(57) Abstract

This invention is an endovascular support device (10) adapted for local delivery of a therapeutic agent (50) for minimizing the rate of restenosis having a cylindrical support body (12) with an inside surface (14), an opposite outside surface (16), and at least one layer of pericardial tissue (20) covering at least a portion of at least a selected one of the inside surface (14), or the outside surface (16) of the cylindrical support body (12). At least one therapeutic agent (50) is disposed on a portion of the support device (10).



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
ΑT	Austria	GE	Georgia	MX	Mexico
ΑÜ	Australia	GN	Guinea	NE	
BB	Barbados	GR	Greece	NE NL	Niger
BE	Belgium	HU	Hungary	NO NO	Netherlands
BF	Burkina Faso	IE	Ireland	NZ	Norway
BG	Bulgaria	IT	Italy	PL	New Zealand
BJ	Benin	JP	Japan	PT	Poland
BR	Brazil .	KE			Portugal
BY	Belarus	KG	Kyrgystan	RO	Romania
CA	Canada	КР	Democratic People's Republic	RU	Russian Federation
CF	Central African Republic		of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SG	Singapore
CI	Côte d'Ivoire	LI	Liechtenstein	SI	Slovenia
CM	Cameroon	ĹK	Sri Lanka	SK	Slovakia
CN	China	LR	Liberia	SN	Senegal
CS	Czechoslovakia	LT		SZ	Swaziland
CZ	Czech Republic	LU	Lithuania	TD	Chad ·
DE	Germany	LV	Luxembourg	TG	Togo
DK	Denmark	MC	Larvia	TJ	Tajikistan
EE	Estonia		Monaco	TT	Trinidad and Tobago
ES	Spain	MD	Republic of Moldova	UA	Ukraine
FI	Finland	MG	Madagascar	UG	Uganda
FR	France	ML	Mali	US	United States of America
GA	Gabon	MN	Mongolia	UZ	Uzbekistan
	00001	MR	Mauritania	VN	Viet Nam

ENDOVASCULAR SUPPORT DEVICE AND METHOD OF USE

5

FIELD OF THE INVENTION

The present invention relates to the field of endovascular support to maintain patency of narrowed vessels and facilitate repair of injured or degenerated vessels by implantation of a device within the target vessel. More specifically, the invention relates to acceptable biological coverings for an endovascular support device and the local delivery of a therapeutic substance into the vascular tissue, as a complementary treatment.

15

20

10

BACKGROUND OF THE INVENTION

Partial and even complete blockage of the vascular system by the formation of an atherosclerotic plaque is a well known and frequent medical problem. Such blockages are often treated with percutaneous transluminal coronary angioplasty (PTCA), also known as balloon angioplasty, or by intravascular stent implantation. PTCA is an alternative to vascular bypass surgery, which includes a surgically exposing incision, and removing, replacing, or bypassing the defective blood vessel. Structures which have previously been used as intraluminal vascular grafts have included coiled stainless steel springs, and grafted stents made out of synthetic material (Dacron or PTFE). Examples

of such prior art devices may be found in U.S. Patent Nos. 5,306,286; 5,026,377; 5,019,085; 5,019,090; 4,913,141; 4,886,062; 4,733,665; and 4,503,569.

In balloon dilatation of vascular stenosis, or blockages, the balloon is inflated within the stenosed vessel, in order to shear and disrupt the wall component of the vessel to obtain an enlarged lumen. This may create a flap or tear at the intima or the media of the vessel wall. The intimal flap can fold down into the lumen and may occlude the vessel. Such occurrences contribute to the high incidence of restinosis which is prevalent. Currently, conventional balloon angioplasty is followed by roughly a 30% to 50% incidence of restenosis.

Vascular prosthetic devices are often utilized in an effort to maintain vessel patency and prevent restinosis. However, vascular prosthetic devices or patches are often associated with increased thrombogenicity of the PTCA site due to the blood contacting the surfaces of the prosthetic device, and result in occlusion of the vesicle. Additionally, synthetic materials used in conventional endovascular prostheses tend to reject coverage by the patient's living tissue, i.e. endothelium, and have collecting surfaces that become thrombogenic sites. Previous biological materials suggested in the art for use in endovascular support have not provided a satisfactory ability to graft to the patient's tissues. Moreover, the body's own repair mechanisms can bring progressive stenotic occlusion due to neointimal fibrosis and hyperplasia. Additionally, an immunological response to foreign material can lead to increased inflammation in response to the prosthetic device.

5

10

15

Systemic therapy aimed at preventing coagulation, a thrombosis locally at the graft site, is often complicated at other sites and produces unwanted, even dangerous side effects. Likewise, systemic treatment with growth mediators or chemotherapeutic agents can produce a hyperplastic or hypoplastic response in tissue not specifically targeted. Existing stent devices, such as dip coated stents, providing locally delivered drugs do not satisfactorily promote grafting or integration of the stent into the patient's endothelium. See for example U.S. Patent Nos. 5,383,928 and 5,102,417.

Therefore, there exists a need in the art for a device and method for supporting endovascular vessels, which provides local therapy for repairing those blood vessels narrowed or occluded by disease, and which provides a biologically acceptable substrate for grafting to the patient. There is a need for such a device which includes providing local therapy resulting in high local concentrations of therapeutic drugs at the treatment site. The art has sought such an expandable intraluminal vascular support graft, and alternatively an independent arterial-vascular fistula, which prevents recurrence of stenosis, to be utilized to support or replace degenerated vein grafts, coronary arteries, and the peripheral arterial and venous system.

However, prior to the development of the present invention, there has been no
expandable intraluminal graft comprised of a biological material which prevents
recurrence of restenosis that can also be utilized to deliver drugs locally to the desired
location.

5

10

SUMMARY OF THE INVENTION

The present invention provides an endovascular support device adapted for local delivery of a therapeutic agent and for minimizing the rate of restinosis. The device has a cylindrical support body having an inside surface and an outside surface. The endovascular support device of the present invention also has at least one layer of pericardial tissue covering at least a portion of the inside surface or the outside surface of the cylindrical support body. The endovascular support device of the present invention is provided with a therapeutic agent disposed on a portion thereof.

10

15

5

BRIEF DESCRIPTIONS OF THE DRAWINGS

Fig. 1 is a perspective view of the endovascular support device in accordance with one embodiment of the invention positioned on a balloon catheter delivery means inside a vesicle.

Fig. 2 is a cross section view of the endovascular support device taken along the plane defined by the line 2-2 in Fig. 1, and positioned within a vesicle.

20

Fig. 3 is a partial cut away view of the pericardial tissue of the endovascular support device with micropores disposed therethrough for delivery of a therapeutic agent.

Fig. 4 is a schematic side view of six (Figs. 4A-4F) alternative embodiments of the endovascular support device of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

Referring to Figures 1-4, the present invention provides an endovascular support device 10 adapted for local delivery of a therapeutic agent 50 and for minimizing the rate of restinosis. The device 10 has a cylindrical support body 12 having an inside surface 14 and an opposite outside surface 16. The cylindrical support body 12 may be constructed of an expandable flexible wire coil or a tubular mesh of multiple coils or rings. Alternatively, the support body 12 may be constructed of a molded polymer, or similarly rigid substance, or a combination of the two.

An expandable support material can permit precise positioning and maintenance of the device 10 within a bodily vesicle 25, with the assistance of an angioplasty balloon 16, as shown in Figure 1. In common use, a guide wire 18 is inserted within the vesicle 25 to the point where endovascular support is desired. An angioplasty balloon 17, or other similar positioning device, carrying the endovascular support device 10 of the present invention, is moved through the vesicle 25 along the guide wire 18. At the desired site, the balloon 17 is inflated and the device 10 is positioned against the walls of the vesicle. The angioplasty balloon 17 and guide wire 18 are then removed, leaving the device 10 in place to support the vesicle 25.

5

The invention contemplates that the endovascular support device 10 can be used within existing and grafted vesicles in a patient. The device 10 may be positioned within a vesicle for a variety of purposes, such as for structural support or to occlude an associated aneurysm, for example. The invention also contemplates that the device 10 can be used independently to provide a vascular replacement for ineffective vesicles, or as a separate fistula as for dialysis.

Referring to Figures 2 and 4, the endovascular support device 10 of the present invention also has at least one layer of pericardial tissue 20 covering at least a portion of the inside surface 14 or the outside surface 16 of the cylindrical support body 12. The pericardial tissue 20 provides a thin-walled membrane made of biological tissue to promote acceptance and fusion with the patient's blood vessel 25 tissue. The support device 10 surface may have an adjustable thickness by varying the number of tissue 20 layers. The pericardium may be chosen from any mammal, but is preferably of porcine, bovine, or human origin.

One skilled in the art will know how to prepare such a pericardial tissue 20 for a homograft or xenograft. For example, porcine pericardium may be retrieved from that surrounding the heart of a sacrificed pig. After shaving the excess fat, the tissue can be immersed in 0.2% glutaraldehyde, which creates permanent strengthening chemical cross-links. The porcine pericardium can then be immersed in porcine albumin solution. When the pericardium is wrapped on the support body 12, it may be adhered to the support body 12 with a biocompatible glue or by careful suturing. The pericardial tissue may be stored in a cold electrolyte solution until used.

5

10

15

Pericardial tissue provides a surprisingly effective biological endovascular support covering. This feature is due, at least in part, to the strength and elasticity of the tissue. The pericardium also has a very low degree of thrombogenicity and is biocompatible with graft recipients. Furthermore, the pericardium serves as an excellent drug delivery means, due to its porosity and high collagen content.

The endovascular support device 10 of the present invention can be provided with at least one therapeutic agent 40 disposed on a portion thereof. By "therapeutic agent" is meant any compound which has a desired pharmacologic effect. By "disposed on" is meant that the therapeutic agent 40 is in contact with at least a portion of the tissue or polymer, described below. For example, the therapeutic agent 40 can be soaked into the tissue or polymer over a period of time, or alternatively, can be injected into a reservoir or cavity created by layers of these materials. The invention contemplates that the device 40 may be made available either presoaked with a therapeutic agent, or provided such that a therapeutic agent is chosen for soaking or injection into the reservoir just prior to placement of the device 10 in the patient.

More specifically, the therapeutic agent 40 can be an anticoagulant, such as D-Phe-Pro-Arg chloromethyl ketone, an RGD peptide-containing compound, heparin, an antithrombin compound, a platelet receptor antagonist, an anti-thrombin antibody, an anti-platelet receptor antibody, aspirin, a prostaglandin inhibitor, a platelet inhibitor or a tick anti-platelet peptide. The therapeutic agent 40 can be a promoter of vascular cell growth, such as a growth factor stimulator, a growth factor receptor agonist, a transcriptional activator, and a translational promoter. Alternatively, the therapeutic

5

10

15

agent 40 can be an inhibitor of vascular cell growth, such as a growth factor inhibitor, a growth factor receptor antagonist, a transcriptional repressor, a translational repressor, an antisense DNA, an antisense RNA, a replication inhibitor, an inhibitory antibody, an antibody directed against growth factors, a bifunctional molecule consisting of a growth factor and a cytotoxin, or a bifunctional molecule consisting of an antibody and a cytotoxin.

The therapeutic agent 40 can be a cholesterol-lowering agent, a vasodilating agent, or other agents which interfere with endogenous vasoactive mechanisms.

Additionally, the therapeutic agent 40 can be a smooth muscle inhibitor, such as: an agent that modulates intracellular calcium binding proteins; a receptor blocker for contractile agonists; an inhibitor of the sodium/hydrogen antiporter; a protease inhibitor; a nitrovasodilator; a phosphodiesterase inhibitor; a phenothiazine; a growth factor receptor agonist; an anti-mitotic agent; an immunosuppressive agent; or a protein kinase inhibitor.

The therapeutic agent 40 may be disposed on all or a portion of the pericardial tissue 20 to utilize the biological properties of the material to absorb different drugs and to release them slowly after deployment to the adjacent tissue. As seen in Figure 3, the pericardial tissue 20 can have a plurality of micropores 22 that extend therethrough for more effective delivery of the therapeutic agent 40. The micropores 22 may also be made to extend only partially into one surface of the tissue 20, and not extend entirely therethrough, such that the therapeutic agent 40 can be directed to diffuse primarily in the direction of the micropores. For example, if only one layer of pericardial tissue 20 is

5

10

15

employed in the endovascular support device 10, the surface of the pericardial tissue 20 intended to face outward can have a plurality of micropores 22 disposed thereon for delivery of the therapeutic agent 50 primarily toward the adjacent walls of the blood vesicle 25, and not toward the interior of the vesicle 25. Micropores 22 can be made in the tissue 20 by a variety of means, including by laser incision.

Alternatively, the therapeutic agent 40 may be disposed on all or a portion of a polymer 50, which can be biodegradable and adapted for slow release of the therapeutic agent 40. A polymer 50 laden with one or more therapeutic agents 40 can be positioned on portion of the cylindrical support body 12, wrapped around a portion of the pericardial tissue 20, or imbedded between multiple layers of pericardial tissues 20. Alternatively, the polymer 50 can be constructed so as to form the cylindrical support body 12 itself, then wrapped at least partially with at least one layer of pericardial tissue 20.

15

20

10

5

A biodegradable polymer 50 such as polylactide, polyanhydride, polyorthoester or polyglycolide, for example can be used. In addition to synthetic polymers, natural polymers can be used, such as amino acid polymers or polysaccharides. The polymer 50 is selected depending on the drug required, the polymer's 50 compatibility with a patient and the ultimate pharmacologic effect desired. For example, if the effect need only last a short period, a thin polymer 50 can be used with a limited amount of drug capable of diffusing from the polymer 50 into the arterial wall or lumen of the vesicle 25.

Alternatively, only the layer closest to the body fluid would contain the therapeutic

agent 40. Another alternative would be to use a polymer 50 which is biodegradable over a long period of time. Naturally, the opposite characteristics would be selected for a desired prolonged release.

The characteristics of the particular polymer 50 for these purposes is well known to the skilled artisan or can be determined by reference to standard references, e.g., Biodegradable Polymers as Drug Delivery Systems, R. Langer and M. Chasin, Eds., Marcel Dekker Inc., New York, NY, USA (1990); Engleberg and Kohn, "Physicomechanical properties of degradable polymers used in medical applications: a comparative study," Biomaterials 12:292-304 (1991); Controlled Release Delivery Systems, T. J. Roseman and S. D. Mansdorf, Eds., Marcel Dekker Inc., New York, NY, USA (1983); and "Controlled Release Technology, Pharmaceutical Applications, ACS Symposium Series, Vol. 348, P. I. Lee and W. R. Good, Eds., American Chemical Society, Washington, D.C., USA (1987).

15

20

Generally, the polymer 50 has a therapeutic agent 40 release rate of between about .001 µg/cm²-min and about 100 µg/cm²-min, especially between about .01 µg/cm²-min and 10 µg/cm²-min. In addition, the polymer 50 generally has a thickness of between about .01 mm and 10 mm, especially between about .1 mm and 1 mm. As can be appreciated, the device 10 can be comprised of two or more different therapeutic agents 40 or two or more different polymers 50 to obtain a desired effect and release rate. In addition, the polymers 50 can have different solubilities or diffusion characteristics to accomplish non-uniform therapeutic agent 40 release.

Figures 4A-4F show a variety of combinations of the device 10 elements: the support body 12, pericardial tissue 20, and polymer 50, contemplated as different embodiments of the present invention. It is understood from the above description that the therapeutic agent 40 may be disposed on any or all of these elements. As discussed above, the therapeutic agent 40 can be, for example, soaked into the pericardial tissue or polymer, laminated on the pericardial tissue or polymer, or injected into a reservoir formed by these materials. Figure 4A shows one embodiment in which the support body 12 has a pericardial tissue 20 disposed on both the inside surface 14 and outside surface 16 thereof. Certainly, the invention contemplates that pericardium tissue 20 can be disposed on either surface alone. Figure 4B shows the support body 12 covered on the inside surface 14 by a layer of pericardial tissue 20, and covered on the outside surface 16 by a layer of polymer 50. Figure 4C shows an embodiment in which the support body 12 is covered on the inside surface 14 with a layer of polymer 50 which is in turn covered by a layer of pericardial tissue 20, and the outside surface 16 of the support body 12 is covered by a layer of polymer 50.

Figure 4D is the embodiment also shown in cross section in Figure 2. This embodiment has the support body 12 sandwiched on the inside surface 14 and the outside surface 16 between two layers of polymer 50, which is in turn sandwiched between two layers of pericardial tissue 20. Figure 4E presents an embodiment wherein the support body 12 is covered on the inside surface 14 by a layer of polymer 50, and on the outside surface 16 by a layer of pericardial tissue 20. Finally, Figure 4F shows an embodiment having a layer of polymer 50 disposed on the inside surface 14 and a layer

10

15

of polymer 50 disposed on the outside surface, which is also coated with a layer of pericardial tissue 20.

The present invention also provides methods of using the device 10 to support an endovascular vessel 25 and to locally deliver a therapeutic agent 40 to minimize the rate of restenosis. More particularly, the invention provides methods for treating or preventing a condition associated with coagulation, thrombus formation, fibrosis and restenosis at treated vascular sites.

Throughout this Specification, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which the invention pertains.

The preceding examples are intended to illustrate, but not limit the invention.

While they are typical of those that might be used, other procedures known to those skilled in the art may be alternatively employed. As used herein, "a" means one or more than one, depending upon the context.

WHAT IS CLAIMED IS:

1. An endovascular support device adapted for local delivery of a therapeutic agent and for minimizing the rate of restenosis, comprising:

- a. a cylindrical support body having an inside surface and an opposite outside surface;
- b. at least one layer of pericardial tissue covering at least a portion of at least a selected one of said inside surface or said outside surface of said cylindrical support body; and,
- c. a therapeutic agent disposed on a portion of said pericardial tissue, thereby adapting said endovascular support device for local delivery of the therapeutic agent and for minimizing the rate of restenosis.
- 2. The endovascular support device of claim 1, wherein said support body is comprised of material selected from the group consisting of a metal and a plastic polymer.
- 3. The endovascular support device of claim 1, further comprising at least one layer of a polymer on at least a portion of said cylindrical support body.
- 4. The endovascular support device of claim 3, wherein a second therapeutic agent is disposed on at least a portion of said polymer.
- 5. The endovascular support device of claim 1, wherein said pericardial tissue is

selected from the group consisting of porcine, bovine and human pericardial tissue.

6. The endovascular support device of claim 1, wherein the pericardial tissue has a plurality of micropores therethrough for delivery of said therapeutic agent.

- 7. The endovascular support device of claim 1, wherein a surface of the pericardial tissue has a plurality of micropores thereon for delivery of said therapeutic agent.
- 8. The endovascular support device of claim 1, wherein said therapeutic agent is disposed upon said pericardial tissue covering at least a portion of said inside surface of said cylindrical support body.
- 9. The endovascular support device of claim 1, wherein said therapeutic agent is an anticoagulant selected from the group consisting of D-Phe-Pro-Arg chloromethyl ketone, an RGD peptide-containing compound, heparin, an antithrombin compound, a platelet receptor antagonist, an anti-thrombin antibody, an anti-platelet receptor antibody, aspirin, a prostaglandin inhibitor, a platelet inhibitor and a tick anti-platelet peptide, and combinations thereof.
- 10. The endovascular support device of claim 1, wherein said therapeutic agent is a promoter of vascular cell growth selected from the group consisting of a growth cor stimulator, a growth factor receptor agonist, a transcriptional activator, and a translational promoter, and combinations thereof.

11. The endovascular support device of claim 1, wherein said therapeutic agent is an inhibitor of vascular cell growth selected from the group consisting of a growth factor inhibitor, a growth factor receptor antagonist, a transcriptional repressor, a translational repressor, an antisense DNA, an antisense RNA, a replication inhibitor, an inhibitory antibody, an antibody directed against growth factors, a bifunctional molecule consisting of a growth factor and a cytotoxin, and a bifunctional molecule consisting of an antibody and a cytotoxin, and combinations thereof.

- 12. The endovascular support device of claim 1, wherein said therapeutic agent is selected from the group consisting of a cholesterol-lowering agent, a vasodilating agent, and agents which interfere with endogenous vasoactive mechanisms, and combinations thereof.
- 13. The endovascular support device of claim 1, wherein said therapeutic agent is a smooth muscle inhibitor selected from the group consisting of an agent that modulates intracellular calcium binding proteins, a receptor blocker for contractile agonists, an inhibitor of the sodium/hydrogen antiporter, a protease inhibitor, a nitrovasodilator, a phosphodiesterase inhibitor, a phenothiazine, a growth factor receptor agonist, an antimitotic agent, an immunosuppressive agent, and a protein kinase inhibitor, and combinations thereof.
- 14. An endovascular support device adapted for local delivery of a therapeutic agent and for minimizing the rate of restenosis, comprising:
 - a. a cylindrical support body having an inside surface and an opposite

outside surface;

b. at least one layer of pericardial tissue covering at least a first portion of at least a selected one of said inside surface or said outside surface of said cylindrical support body;

- c. a polymer covering at least a second portion of at least a selected one of said inside surface or said outside surface of said cylindrical support body; and
- d. a therapeutic agent disposed on said polymer, thereby adapting said endovascular support device for local delivery of the therapeutic agent and for minimizing the rate of restenosis.
- 15. The endovascular support device of claim 14, wherein at least a portion of said pericardial tissue is disposed on at least a portion of said polymer.
- 16. The endovascular support device of claim 14, wherein at least a portion of said pericardial tissue is disposed between at least a portion of said cylindrical support body and at least a portion of said polymer.
- 17. The endovascular support device of claim 14, wherein a second therapeutic agent is disposed on at least a portion of said pericardial tissue.
- 18. The endovascular support device of claim 14, wherein said pericardial tissue is selected from the group consisting of porcine, bovine and human pericardial tissue.
- 19. The endovascular support device of claim 14, wherein the pericardial tissue has a

plurality of micropores therethrough for delivery of said therapeutic agent.

20. The endovascular support device of claim 14, wherein a surface of the pericardial tissue has a plurality of micropores thereon for delivery of said therapeutic agent.

- 21. The endovascular support device of claim 14, wherein said therapeutic agent is an anticoagulant selected from the group consisting of D-Phe-Pro-Arg chloromethyl ketone, an RGD peptide-containing compound, heparin, an antithrombin compound, a platelet receptor antagonist, an anti-thrombin antibody, an anti-platelet receptor antibody, aspirin, a prostaglandin inhibitor, a platelet inhibitor and a tick anti-platelet peptide, and combinations thereof.
- 22. The endovascular support device of claim 14, wherein said therapeutic agent is a promoter of vascular cell growth selected from the group consisting of a growth factor stimulator, a growth factor receptor agonist, a transcriptional activator, and a translational promoter, and combinations thereof.
- 23. The endovascular support device of claim 14, wherein said therapeutic agent is an inhibitor of vascular cell growth selected from the group consisting of a growth factor inhibitor, a growth factor receptor antagonist, a transcriptional repressor, a translational repressor, an antisense DNA, an antisense RNA, a replication inhibitor, an inhibitory antibody, an antibody directed against growth factors, a bifunctional molecule consisting of a growth factor and a cytotoxin, and a bifunctional molecule consisting of

an antibody and a cytotoxin, and combinations thereof.

24. The endovascular support device of claim 14, wherein said therapeutic agent is selected from the group consisting of a cholesterol-lowering agent, a vasodilating agent, and agents which interfere with endogenous vasoactive mechanisms, and combinations thereof.

- 25. The endovascular support device of claim 14, wherein said therapeutic agent is a smooth muscle inhibitor selected from the group consisting of an agent that modulates intracellular calcium binding proteins, a receptor blocker for contractile agonists, an inhibitor of the sodium/hydrogen antiporter, a protease inhibitor, a nitrovasodilator, a phosphodiesterase inhibitor, a phenothiazine, a growth factor receptor agonist, an antimitotic agent, an immunosuppressive agent, and a protein kinase inhibitor, and combinations thereof.
- 26. An endovascular support device adapted for local delivery of a therapeutic agent and for minimizing the rate of restenosis, comprising:
 - a. a cylindrical support body having an inside surface and an opposite outside surface;
 - b. a first pericardial tissue covering at least a first portion of at least a selected one of said inside surface or said outside surface of said cylindrical support body; and
 - c. a second pericardial tissue covering at least a second portion of at least a selected one of said inside surface or said outside surface of said cylindrical

support body; and


d. at least one therapeutic agent disposed in a reservoir between said first pericardial tissue and said second pericardial tissue, thereby adapting said endovascular support device for local delivery of the therapeutic agent and for minimizing the rate of restenosis.


- 27. The endovascular support device of claim 26, further comprising a polymer on at least a portion of said cylindrical support body.
- 28. The endovascular support device of claim 26, wherein a second therapeutic agent is disposed on at least a portion of said polymer.
- 29. The endovascular support device of claim 26, wherein said pericardial tissue is selected from the group consisting of porcine, bovine and human pericardial tissue.
- 30. The endovascular support device of claim 26, wherein the pericardial tissue has a plurality of micropores therethrough for delivery of said therapeutic agent.
- 31. The endovascular support device of claim 26, wherein a surface of the pericardial tissue has a plurality of micropores thereon for delivery of said therapeutic agent.
- 32. A method of supporting an endovascular vessel and locally delivering a therapeutic agent and minimizing the rate of restenosis, comprising inserting in the

vessel an endovascular support device comprising:

a. a cylindrical support body having an inside surface and an opposite outside surface;

- b. at least one layer of pericardial tissue covering at least a portion of at least a selected one of said inside surface or said outside surface of said cylindrical support body; and,
- c. a therapeutic agent disposed on at least a portion of said pericardial tissue, thereby supporting the endovascular vessel and locally delivering the therapeutic agent and minimizing the rate of restenosis.
- 33. A method of replacing a vascular vessel and locally delivering a therapeutic agent and minimizing the rate of restenosis, comprising replacing the vessel with an endovascular support device comprising:
 - a. a cylindrical support body having an inside surface and an opposite outside surface;
 - b. at least one layer of pericardial tissue covering at least a portion of at least a selected one of said inside surface or said outside surface of said cylindrical support body; and,
 - c. a therapeutic agent disposed on at least a portion of said pericardial tissue, thereby replacing the endovascular vessel and locally delivering the therapeutic agent and minimizing the rate of restenosis.





INTERNATIONAL SEARCH REPORT

International application No. PCT/US96/13907

		PC170390/1390/			
A. CL	ASSIFICATION OF SUBJECT MATTER				
IPC(6)	:A61F 2/04, 06, 24 54				
According	:42A/423; 600/36; 623/1, 2, 11, 12, 66 to International Patent Classification (IPC) or to both national classification are				
B. FIE	CLDS SEARCHED	nd IPC			
	documentation scarched (classification system followed by classification symbol				
U.S. :	424/423; 600/36; 623/1, 2, 11, 12, 66	ols)			
Documents	ation searched other than minimum documentation to the extent that such docume				
		aus are included in the fields searche	×d		
Electronic (data base consulted during the international search (name of data base and, wh	ere practicable seemb to an a			
APS		presentative, search terms used)			
C. DOC	TIMPHTS CONTENTS		•		
	CUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant	passages Relevant to claim			
Y			NO.		
•	US, A, 5,383,927 (De GOICOECHEA ET AL.) 24	January 1-33			
	1995, see column 6 lines 4-20, column 7 lines 18 Fig. 1B.	-65, and			
	1 ig. 15.				
i			ĺ		
	,				
/	US, A, 4,729,139 (NASHEF) 08 March 1988, see				
İ	2 lines 38-62.	e column 1-33	•		
′	US, A, 5,263,992 (GUIRE) 23 November 19	102			
- 1	column 4 lines 24-50.	95, see 4, 17, 28			
, 1					
	US, A, 5,399,352 (HANSON) 21 March 1995, see	column 9-13, 21-	25		
ļ	5 line 59 to column 6 line 9, and Figs. 1, 2, 5 and	7. 32, 33	25,		
	•		1		
		1 .			
			l		
7					
	er documents are listed in the continuation of Box C. See patent fan	nily annex.	\Box		
	cial energories of cited documents:	shed after the interactional filing date or prior			
	e part of paracular relevance or the art which m not considered principle or theory a	ict with the application but cited to understand underlying the invention	•••		
certi	are document published on or after the interestional filing date. "X" document of particu	the relevance the allies at the st			
docu	MERCENT Which many throw doubts on priority claim(a) or which is whom the document	CONTRACTO CONTRACTOR IN TRACTOR OF THE PROPERTY OF	 >		
•	document of particu	the relevance; the claimed invention cannot			
Dica:	as only declosure, tax, exhibition or other combined with one of	we an inventive step when the document we more other such documents, such combinate cross skilled in the art.			
docu	sment published prior to the international Clina day but to	erson skilled in the art f the mene patent family			
	consisting of the consistency of				
04 OCTOBER 1996 Date of mailing of the international search report 24 OCTOBER 1996					
~ OCTOB	74 UC 1	YYb	- 1		
ame and ma	ailing address of the ISA/US Authorized officer				
Box PCT	er of Palents and Trademarks	ele Kolina	•••		
Washington, crimile No.	D.C. 20231 FRANCIS K. CUDDI	THY	1		
m PCT/IC	. (703) 305-3590 Telephone No. (703) 3	308-2996	- 1		

Date: 15/11/2001

| JOHN O'BRIEN & ASSOCIATES Attn. O'Brien, John A. 3rd Floor, Duncairn House 14 Carysfort Avenue Blackrock, County Dublin IRELAND